

Chromatic Roots of a Ring of Four Cliques

F.M. Dong*

Mathematics and Mathematics Education, National Institute of Education
Nanyang Technological University, Singapore

Gordon Royle

School of Mathematics & Statistics, University of Western Australia
Email: gordon@maths.uwa.edu.au

Dave Wagner

Department of Combinatorics and Optimization, University of Waterloo
Email: dgwagner@math.uwaterloo.ca

April 27, 2011

Abstract

For any positive integers a, b, c, d , let $R_{a,b,c,d}$ be the graph obtained from the complete graphs K_a, K_b, K_c and K_d by adding edges joining every vertex in K_a and K_c to every vertex in K_b and K_d . This paper shows that for arbitrary positive integers a, b, c and d , every root of the chromatic polynomial of $R_{a,b,c,d}$ is either a real number or a complex number with its real part equal to $(a + b + c + d - 1)/2$.

Keywords: graph, chromatic polynomial, chromatic root, ring of cliques

1 Introduction

A *ring of cliques* is a graph whose vertex set is the disjoint union of cliques, arranged in a cyclic order, such that the vertices of each clique are joined to all the vertices in the two neighbouring cliques. If the cliques have size a_1, a_2, \dots, a_n then we denote this graph by R_{a_1, a_2, \dots, a_n} . Figure 1 shows the graph $R_{2,2,3,3}$.

Graphs with this structure have occurred several times previously in the study of chromatic polynomials and their roots. In particular, in proving that there are non-chordal graphs with integer chromatic roots, Read [6] considered the graphs in this family with $a_1 = 1$ (and he also used slightly different notation). Rings of cliques cropped up again recently in a preliminary investigation of the *algebraic properties* of chromatic roots (Cameron [1]) and in the course

*Corresponding author. Email: fengming.dong@nie.edu.sg.




Figure 1: The graph $R_{2,2,3,3}$

of this investigation, the chromatic roots of many of these graphs were computed. When the chromatic roots of the ring-of-clique graphs with *exactly four cliques* and a fixed number of vertices were plotted, an intriguing pattern was observed — all the complex chromatic roots lie on a single vertical line. Figure 2 shows the union of the chromatic roots of the 12-vertex graphs of the form $R_{a,b,c,d}$.

Faced with such a striking empirically-observed pattern, we were led to explain it theoretically. This appears to require a surprisingly intricate argument, but eventually we obtain the following result:

Theorem 1 *For arbitrary non-negative integers a, b, c and d the chromatic roots of $R_{a,b,c,d}$ are either real, or complex with real part equal to $(a + b + c + d - 1)/2$.*

The overall structure of the argument is as follows: the chromatic polynomial $P(R_{a,b,c,d}, \lambda)$ is first expressed as the product of linear factors and a factor $Q_{a,b,c,d}(\lambda)$. It then suffices to show that the complex roots of $Q_{a,b,c,d}(\lambda)$ all lie on the vertical line $\Re(\lambda) = (a + b + c + d - 1)/2$ in the complex λ -plane. Next the polynomial $F_{a,p,q,n}(z)$ is defined to be $Q_{a,b,c,d}(z + (a + b + c + d - 1)/2)$ thus translating the vertical line supposed to contain the roots to the imaginary axis and also reparameterizing the problem (in a somewhat counterintuitive way). Then $F_{a,p,q,n}$ is shown to be an *even* polynomial and we define a fourth polynomial $W_{a,p,q,n}$ by $W_{a,p,q,n}(z^2) = F_{a,p,q,n}(z)$. The proof is completed by demonstrating that $W_{a,p,q,n}$ is real-rooted using polynomial interleaving techniques, and therefore $F_{a,p,q,n}$ has only real or pure imaginary roots as required.




Figure 2: Chromatic roots of the graphs $R_{a,b,c,d}$ where $a + b + c + d = 12$.

2 Basics

For any graph G and any positive integer λ , let $P(G, \lambda)$ be the number of mappings ϕ from $V(G)$ to $\{1, 2, \dots, \lambda\}$ such that $\phi(u) \neq \phi(v)$ for every two adjacent vertices u and v in G . It is well-known that $P(G, \lambda)$ is a polynomial in λ , called *the chromatic polynomial* of G .

The chromatic polynomial of a graph G has the following properties (see, for instance, [3, 5, 7, 8]), which we will apply later.

Proposition 1 *Let G be a simple graph of order at least 2.*

(i) *If u and v are two non-adjacent vertices in G , then*

$$P(G, \lambda) = P(G + uv, \lambda) + P(G/uv, \lambda), \quad (1)$$

where $G + uv$ is the graph obtained from G by adding the edge joining u and v , and G/uv is the graph obtained from G by identifying u and v and removing all parallel edges but one.

(ii) *If u is a vertex in G which is adjacent to all other vertices in G , then*

$$P(G, \lambda) = \lambda P(G - u, \lambda - 1), \quad (2)$$

where $G - u$ is the graph obtained from G by removing u .

If $a = 0$, $R_{a,b,c,d}$ is a chordal graph and its chromatic polynomial is

$$P(R_{0,b,c,d}, \lambda) = \frac{(\lambda)_{b+c}(\lambda)_{c+d}}{(\lambda)_c}, \quad (3)$$

and if $a \geq 1$ and $c \geq 1$, then applying Proposition 1 repeatedly yields that

$$P(R_{a,b,c,d}, \lambda) = \lambda P(R_{a-1,b,c,d}, \lambda - 1) + c\lambda P(R_{a-1,b,c-1,d}, \lambda - 1). \quad (4)$$

For a non-negative integer a and real numbers b, c and d , define a polynomial $Q_{a,b,c,d}(z)$ in z as follows: $Q_{0,b,c,d}(z) = 1$ and for $a \geq 1$,

$$Q_{a,b,c,d}(z) = (z - b - c)(z - c - d)Q_{a-1,b,c,d}(z - 1) + c(z - a - c + 1)Q_{a-1,b,c-1,d}(z - 1). \quad (5)$$

It is clear that $Q_{a,b,c,d}(z)$ is a polynomial of order $2a$ in z .

Proposition 2 *Let a, b, c and d be any non-negative integers. Then*

$$P(R_{a,b,c,d}, \lambda) = \frac{(\lambda)_{b+c}(\lambda)_{c+d}}{(\lambda)_{a+c}} Q_{a,b,c,d}(\lambda). \quad (6)$$

Proof. If $a = 0$, then (6) follows from (3) and the definition of $Q_{a,b,c,d}(\lambda)$. Now assume that $a \geq 1$. By (4) and induction, we have

$$\begin{aligned} P(R_{a,b,c,d}, \lambda) &= \lambda P(R_{a-1,b,c,d}, \lambda - 1) + c\lambda P(R_{a-1,b,c-1,d}, \lambda - 1) \\ &= \lambda \frac{(\lambda - 1)_{b+c}(\lambda - 1)_{c+d}}{(\lambda - 1)_{a+c-1}} Q_{a-1,b,c,d}(\lambda - 1) \\ &\quad + c\lambda \frac{(\lambda - 1)_{b+c-1}(\lambda - 1)_{c+d-1}}{(\lambda - 1)_{a+c-2}} Q_{a-1,b,c-1,d}(\lambda - 1) \\ &= \frac{(\lambda)_{b+c}(\lambda)_{c+d}}{(\lambda)_{a+c}} [(\lambda - b - c)(\lambda - c - d)Q_{a-1,b,c,d}(\lambda - 1) \\ &\quad + c(\lambda - a - c + 1)Q_{a-1,b,c-1,d}(\lambda - 1)]. \end{aligned} \quad (7)$$

The result then follows. □

Assume that $\binom{x}{0} = 1$ and $\binom{x}{r} = x(x-1)\cdots(x-r+1)/r!$ for any positive integer r and any complex number x .

Proposition 3 *For any non-negative integer a and real numbers b, c and d ,*

$$Q_{a,b,c,d}(\lambda) = a! \sum_{i=0}^a i!(a-i)! \binom{c}{i} \binom{\lambda - b - c}{a-i} \binom{\lambda - c - d}{a-i} \binom{\lambda - a - c + i}{i}. \quad (8)$$

Proof. It is trivial if $a = 0$ as $Q_{0,b,c,d}(z) = 1$. Now assume that $a \geq 1$. By (5) and induction,

$$\begin{aligned}
& Q_{a,b,c,d}(\lambda) \\
&= (\lambda - b - c)(\lambda - c - d)Q_{a-1,b,c,d}(\lambda - 1) + c(\lambda - a - c + 1)Q_{a-1,b,c-1,d}(\lambda - 1) \\
&= (\lambda - b - c)(\lambda - c - d)(a - 1)! \sum_{i=0}^{a-1} \left\{ i!(a - i - 1)! \binom{c}{i} \binom{\lambda - b - c - 1}{a - i - 1} \right. \\
&\quad \left. \binom{\lambda - c - d - 1}{a - i - 1} \binom{\lambda - a - c + i}{i} \right\} \\
&\quad + c(\lambda - a - c + 1)(a - 1)! \sum_{i=0}^{a-1} \left\{ i!(a - i - 1)! \binom{c - 1}{i} \binom{\lambda - b - c}{a - i - 1} \right. \\
&\quad \left. \binom{\lambda - c - d}{a - i - 1} \binom{\lambda - a - c + i + 1}{i} \right\} \\
&= (a - 1)! \sum_{i=0}^{a-1} i!(a - i)!(a - i) \binom{c}{i} \binom{\lambda - b - c}{a - i} \binom{\lambda - c - d}{a - i} \binom{\lambda - a - c + i}{i} \\
&\quad + (a - 1)! \sum_{i=0}^{a-1} i!(a - i - 1)!(i + 1)^2 \binom{c}{i + 1} \binom{\lambda - b - c}{a - i - 1} \binom{\lambda - c - d}{a - i - 1} \binom{\lambda - a - c + i + 1}{i + 1} \\
&= (a - 1)! \sum_{i=0}^{a-1} i!(a - i)!(a - i) \binom{c}{i} \binom{\lambda - b - c}{a - i} \binom{\lambda - c - d}{a - i} \binom{\lambda - a - c + i}{i} \\
&\quad + (a - 1)! \sum_{i=1}^a (i - 1)!(a - i)! i^2 \binom{c}{i} \binom{\lambda - b - c}{a - i} \binom{\lambda - c - d}{a - i} \binom{\lambda - a - c + i}{i} \\
&= a! \sum_{i=0}^a i!(a - i)! \binom{c}{i} \binom{\lambda - b - c}{a - i} \binom{\lambda - c - d}{a - i} \binom{\lambda - a - c + i}{i}. \tag{9}
\end{aligned}$$

The result then follows. \square

For any non-negative integer a and real numbers p, q, n , define

$$F_{a,p,q,n}(z) = a! \sum_{i=0}^a i!(a - i)! \binom{a + p + q - 1}{i} \binom{z + n + i - 1}{i} \binom{z - p}{a - i} \binom{z - q}{a - i}. \tag{10}$$

Then (8) and (10) implies that $Q_{a,b,c,d}(z + (a + b + c + d - 1)/2) = F_{a,p,q,n}(z)$, where

$$\begin{cases} p = (b + c - a - d + 1)/2 \\ q = (c + d - a - b + 1)/2 \\ n = (b + d - a - c + 1)/2. \end{cases} \tag{11}$$

In the next section, we shall show that $F_{a,p,q,n}(z)$ is an even polynomial in z , and the polynomial obtained from $F_{a,p,q,n}(z)$ by replacing z^2 by z (i.e., $W_{a,p,q,n}(z)$ defined on Page 9) has only real roots for an arbitrary positive integer a and arbitrary real numbers p, q, n satisfying the condition that $p + q, p + n$ and $q + n$ are all non-negative (see Proposition 10). This result implies that every root of $F_{a,p,q,n}(z)$ is either a real number or a complex number with its real part equal to 0 if a is a positive integer and $p + q, p + n$ and $q + n$ are all non-negative real

numbers. For arbitrary positive integers a, b, c, d , if $a \leq \min\{b, c, d\}$ and p, q and n are given in (11), then $p + q = c - a + 1 > 0$, $p + n = b - a + 1 > 0$ and $q + n = d - a + 1 > 0$. Since $Q_{a,b,c,d}(z + (a + b + c + d - 1)/2) = F_{a,p,q,n}(z)$, where p, q and n are given in (11), the following result is obtained.

Proposition 4 *For arbitrary positive integers a, b, c and d , if $a \leq \min\{b, c, d\}$, then every root of $Q_{a,b,c,d}(z)$ is either a real number or a complex number with its real part equal to $(a + b + c + d - 1)/2$. Therefore, for arbitrary non-negative integers a, b, c and d , every root of $P(R_{a,b,c,d}, \lambda)$ is either a real number or a complex number with its real part equal to $(a + b + c + d - 1)/2$. \square*

3 The polynomial $F_{a,p,q,n}(z)$

From the definition of $F_{a,p,q,n}(z)$, we have $F_{0,p,q,n}(z) = 1$ and $F_{1,p,q,n}(z) = z^2 + pq + pn + qn$. We shall show that $F_{a,p,q,n}(z)$ has a recursive expression in terms of $F_{a-1,p,q,n}(z)$ and $F_{a-2,p,q,n}(z)$. We first prove two properties of $F_{a,p,q,n}(z)$.

Proposition 5 *For any integer $a \geq 1$ and arbitrary real numbers p, q, n , if $p + q = 0$, then*

$$F_{a,p,q,n}(z) = (z - p)(z - q)F_{a-1,p+1,q+1,n}(z). \quad (12)$$

Proof. For $a \geq 1$,

$$\begin{aligned} F_{a,p,q,n}(z) &= a! \sum_{i=0}^a i!(a-i)! \binom{a-1}{i} \binom{z+n+i-1}{i} \binom{z-p}{a-i} \binom{z-q}{a-i} \\ &= (z-p)(z-q)(a-1)! \\ &\quad \sum_{i=0}^{a-1} i!(a-1-i)! \binom{a}{i} \binom{z+n+i-1}{i} \binom{z-p-1}{a-1-i} \binom{z-q-1}{a-1-i} \\ &= (z-p)(z-q) \times F_{a-1,p+1,q+1,n}(z). \end{aligned} \quad (13)$$

\square

Proposition 6 *For any integer $a \geq 1$ and arbitrary real numbers p, q, n ,*

$$F_{a,p+1,q,n}(z) - F_{a,p,q,n}(z) = a(a+n+q-1)F_{a-1,p+1,q,n}(z). \quad (14)$$

Proof. For $a \geq 1$,

$$\begin{aligned} &F_{a,p+1,q,n}(z) - F_{a,p,q,n}(z) \\ &= a! \sum_{i=0}^a \left\{ i!(a-i)! \binom{z+n+i-1}{i} \binom{z-q}{a-i} \right. \end{aligned}$$

$$\begin{aligned}
& \left(\binom{a+p+q}{i} \binom{z-p-1}{a-i} - \binom{a+p+q-1}{i} \binom{z-p}{a-i} \right) \Big\} \\
= & a! \sum_{i=0}^a \left\{ i!(a-i)! \binom{z+n+i-1}{i} \binom{z-q}{a-i} \right. \\
& \left. \left(\binom{a+p+q-1}{i-1} \binom{z-p-1}{a-i} - \binom{a+p+q-1}{i} \binom{z-p-1}{a-i-1} \right) \right\} \\
= & a! \sum_{i=1}^a i!(a-i)! \binom{z+n+i-1}{i} \binom{z-q}{a-i} \binom{a+p+q-1}{i-1} \binom{z-p-1}{a-i} \\
& - a! \sum_{i=0}^{a-1} i!(a-i)! \binom{z+n+i-1}{i} \binom{z-q}{a-i} \binom{a+p+q-1}{i} \binom{z-p-1}{a-i-1} \\
= & a! \sum_{i=0}^{a-1} (i+1)!(a-i-1)! \binom{z+n+i}{i+1} \binom{z-q}{a-i-1} \binom{a+p+q-1}{i} \binom{z-p-1}{a-i-1} \\
& - a! \sum_{i=0}^{a-1} i!(a-i)! \binom{z+n+i-1}{i} \binom{z-q}{a-i} \binom{a+p+q-1}{i} \binom{z-p-1}{a-i-1} \\
= & (n+q+a-1)a! \\
& \sum_{i=0}^{a-1} i!(a-1-i)! \binom{z+n+i-1}{i} \binom{z-q}{a-1-i} \binom{a+p+q-1}{i} \binom{z-p-1}{a-i-1} \\
= & a(a+n+q-1)F_{a-1,p+1,q,n}(z). \tag{15}
\end{aligned}$$

□

Now we can prove that $F_{a,p,q,n}(z)$ can be expressed in terms of $F_{a-1,p,q,n}(z)$ and $F_{a-2,p,q,n}(z)$.

Proposition 7 *Let p, q, n be arbitrary real numbers. For any integer $a \geq 2$,*

$$\begin{aligned}
F_{a,p,q,n}(z) = & (z^2 + (a-1)(2p+2q+2n+2a-3) + pq + pn + qn)F_{a-1,p,q,n}(z) \\
& - (a-1)(p+q+a-2)(q+n+a-2)(p+n+a-2)F_{a-2,p,q,n}(z). \tag{16}
\end{aligned}$$

Proof. By the definition of $F_{a,p,q,n}(z)$, we have $F_{0,p,q,n}(z) = 1$, $F_{1,p,q,n}(z) = z^2 + pq + pn + qn$ and

$$\begin{aligned}
F_{2,p,q,n}(z) = & z^4 + (2q + 2pq + 1 + 2pn + 2p + 2qn + 2n)z^2 + pq^2 + pq \\
& + qn + q^2n + p^2q^2 + p^2n^2 + p^2q + 4pqn + pn^2 + 2p^2qn \\
& + pn + 2pq^2n + 2pqn^2 + qn^2 + q^2n^2 + p^2n. \tag{17}
\end{aligned}$$

Thus it can be verified that (16) holds when $a = 2$.

Assume that (16) holds for every integer $2 \leq a < k$, where $k \geq 3$. Now consider the case that $a = k$.

By the definition of $F_{a,p,q,n}(z)$, $F_{a,p,q,n}(z)$ is also a polynomial of order a in p . Let q, n, z be any fixed real numbers. If (16) holds for all numbers p in the set $\{-q+r : r = 0, 1, 2, \dots\}$, then the result is proven.

By assumption on a , (16) holds for $F_{a-1,-q+1,q+1,n}(z)$ and thus

$$\begin{aligned} & F_{a-1,-q+1,q+1,n}(z) \\ = & (z^2 - 5a + 2an + 2a^2 + 3 - 2n - q^2)F_{a-2,-q+1,q+1,n}(z) \\ & - (a-2)(a-1)(-q-2+n+a)(q-2+n+a)F_{a-3,-q+1,q+1,n}(z). \end{aligned}$$

By Proposition 5, for any integer $m \geq 1$,

$$F_{m,-q,q,n}(z) = (z^2 - q^2)F_{m-1,-q+1,q+1,n}(z).$$

Hence

$$\begin{aligned} F_{a,-q,q,n}(z) &= (z^2 - 5a + 2an + 2a^2 + 3 - 2n - q^2)F_{a-1,-q,q,n}(z) \\ &\quad - (a-2)(a-1)(-q-2+n+a)(q-2+n+a)F_{a-2,-q,q,n}(z), \end{aligned}$$

implying that (16) holds for $F_{a,-q,q,n}(z)$.

In the remaining part of this proof, we shall show that if (16) holds for $F_{a,p,q,n}(z)$, then (16) holds for $F_{a,p+1,q,n}(z)$. Assume (16) holds for $F_{a,p,q,n}(z)$, and so

$$\begin{aligned} F_{a,p,q,n}(z) &= (z^2 + (a-1)(2p+2q+2n+2a-3) + pq + pn + qn)F_{a-1,p,q,n}(z) \\ &\quad - (a-1)(p+q+a-2)(q+n+a-2)(p+n+a-2)F_{a-2,p,q,n}(z). \end{aligned} \quad (18)$$

By assumption on a , (16) holds for $F_{a-1,p+1,q,n}(z)$ and so

$$\begin{aligned} & F_{a-1,p+1,q,n}(z) \\ = & (z^2 + (a-2)(2p+2q+2n+2a-3) + (p+1)(n+q) + qn)F_{a-2,p+1,q,n}(z) \\ & - (a-2)(p+q+a-2)(q+n+a-3)(p+n+a-2)F_{a-3,p+1,q,n}(z). \end{aligned} \quad (19)$$

By Proposition 6, (18) and (19), we have

$$\begin{aligned} & F_{a,p+1,q,n}(z) \\ = & F_{a,p,q,n}(z) + a(a+q+n-1)F_{a-1,p+1,q,n}(z) \\ = & (z^2 + (a-1)(2p+2q+2n+2a-3) + pq + pn + qn)F_{a-1,p,q,n}(z) \\ & - (a-1)(p+q+a-2)(q+n+a-2)(p+n+a-2)F_{a-2,p,q,n}(z) \\ & + a(a+q+n-1)F_{a-1,p+1,q,n}(z) \\ = & (z^2 + (a-1)(2p+2q+2n+2a-3) + pq + pn + qn) \\ & (F_{a-1,p+1,q,n}(z) - (a-1)(a+q+n-2)F_{a-2,p+1,q,n}(z)) \\ & - (a-1)(p+q+a-2)(q+n+a-2)(p+n+a-2) \\ & (F_{a-2,p+1,q,n}(z) - (a-2)(a+q+n-3)F_{a-3,p+1,q,n}(z)) \\ & + a(a+q+n-1)F_{a-1,p+1,q,n}(z) \end{aligned}$$

$$\begin{aligned}
&= (z^2 + (a-1)(2p+2q+2n+2a-3) + pq + pn + qn) \\
&\quad (F_{a-1,p+1,q,n}(z) - (a-1)(a+q+n-2)F_{a-2,p+1,q,n}(z)) \\
&\quad - (a-1)(p+q+a-2)(q+n+a-2)(p+n+a-2)F_{a-2,p+1,q,n}(z) \\
&\quad + (a-1)(a+q+n-2)(-F_{a-1,p+1,q,n}(z) + \\
&\quad (z^2 + (a-2)(2p+2q+2n+2a-3) + (p+1)(n+q) + qn)F_{a-2,p+1,q,n}(z)) \\
&\quad + a(a+q+n-1)F_{a-1,p+1,q,n}(z) \\
&= (z^2 + (a-1)(2p+2q+2n+2a-1) + (p+1)(n+q) + qn)F_{a-1,p+1,q,n}(z) \\
&\quad - (a-1)(p+q+a-1)(q+n+a-2)(p+n+a-1)F_{a-2,p+1,q,n}(z).
\end{aligned}$$

Thus (16) holds for $F_{a,p+1,q,n}(z)$. Hence (16) holds for $F_{a,p,q,n}(z)$ for all numbers p in the set $\{q+r : r=0,1,2,\dots\}$ and therefore the result is proved. \square

Since $F_{0,p,q,n}(z) = 1$ and $F_{1,p,q,n}(z) = z^2 + pq + pn + qn$, Proposition 7 implies that $F_{a,p,q,n}(z)$ is an even polynomial in z . For any non-negative integer a and real numbers p, q, n , let $W_{a,p,q,n}(z)$ be the polynomial in z defined as follows: $W_{0,p,q,n}(z) = 1$, $W_{1,p,q,n}(z) = z + pq + pn + qn$ and for $a \geq 2$,

$$\begin{aligned}
W_{a,p,q,n}(z) &= (z + (a-1)(2p+2q+2n+2a-3) + pq + pn + qn)W_{a-1,p,q,n}(z) \\
&\quad - (a-1)(p+q+a-2)(q+n+a-2)(p+n+a-2)W_{a-2,p,q,n}(z). \quad (20)
\end{aligned}$$

Thus it is clear that $F_{a,p,q,n}(z) = W_{a,p,q,n}(z^2)$.

For two non-increasing sequences (a_1, a_2, \dots, a_m) and (b_1, b_2, \dots, b_n) of real numbers, we say the first *interleaves* the second if $m = n+1$ and $(a_1, b_1, a_2, b_2, \dots, a_n, b_n, a_{n+1})$ is a non-increasing sequence, or $m = n$ and $(a_1, b_1, a_2, b_2, \dots, a_n, b_n)$ is a non-increasing sequence. If both polynomials $f(x)$ and $g(x)$ in x with real coefficients have only real roots and the non-increasing sequence formed by all roots of $f(x)$ interleaves the non-increasing sequence formed by all roots of $g(x)$, then we say $f(x)$ *interleaves* $g(x)$. We need to apply the following result given in Section 1.3 of [4]. More details on polynomials with only real roots can be found in [2, 4].

Proposition 8 ([4]) *Let $f(x)$ and $g(x)$ be polynomials with real coefficients and with positive leading coefficients and u and v be any real numbers. If $f(x)$ interleaves $g(x)$ and $v \leq 0$, then $(x-u)f(x) + vg(x)$ interleaves $f(x)$. \square*

Applying Proposition 8, we can get the following result.

Proposition 9 *Let a be any positive integer and p, q, n be any real numbers.*

- (i) *If $(p+q)(n+q)(n+p) \geq 0$, then $W_{2,p,q,n}(z)$ interleaves $W_{1,p,q,n}(z)$.*

- (ii) If $a \geq 3$, $(p + q + a - 2)(q + n + a - 2)(p + n + a - 2) \geq 0$ and $W_{a-1,p,q,n}(z)$ interleaves $W_{a-2,p,q,n}(z)$, then $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$.

Proof. By the definition of $W_{a,p,q,n}(z)$, $W_{1,p,q,n}(z) = z + pq + pn + qn$ and

$$\begin{aligned} W_{2,p,q,n}(z) &= (z + 2p + 2q + 2n + 1 + pq + pn + qn)(z + pq + pn + qn) \\ &\quad - (p + q)(q + n)(p + n). \end{aligned} \quad (21)$$

As the only root of $W_{1,p,q,n}(z)$ is $-pq - pn - qn$ and $W_{2,p,q,n}(-pq - pn - qn) = -(p + q)(n + q)(n + p) \leq 0$, $W_{2,p,q,n}(z)$ interleaves $W_{1,p,q,n}(z)$. So (i) holds.

By Proposition 7,

$$\begin{aligned} F_{a,p,q,n}(z) &= (z^2 + (a - 1)(2p + 2q + 2n + 2a - 3) + pq + pn + qn)F_{a-1,p,q,n}(z) \\ &\quad - (a - 1)(p + q + a - 2)(q + n + a - 2)(p + n + a - 2)F_{a-2,p,q,n}(z). \end{aligned} \quad (22)$$

Since $-(a - 1)(p + q + a - 2)(q + n + a - 2)(p + n + a - 2) \leq 0$ and $W_{a-1,p,q,n}(z)$ interleaves $W_{a-2,p,q,n}(z)$, Proposition 8 implies that $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$. Hence (ii) holds. \square

Notice that $W_{a,p,q,n}(z) = W_{a,q,p,n}(z) = W_{a,n,q,p}(z)$ holds for arbitrary real numbers p, q, n and non-negative integer a , we assume that $p \leq q \leq n$ in the following.

Proposition 10 *Let p, q, n be arbitrary real numbers with $p \leq q \leq n$ and $p + q \geq 0$. Then, for every integer $a \geq 2$, $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$. Therefore, for every positive integer a , $W_{a,p,q,n}(z)$ has only real roots and every root of $F_{a,p,q,n}(z)$ is either a real number or a complex number with its real part equal to 0.*

Proof. Since $p + q \geq 0$ and $p \leq q \leq n$, we have $q + n \geq p + n \geq 0$ and so Proposition 9 (i) implies that $W_{2,p,q,n}(z)$ interleaves $W_{1,p,q,n}(z)$. Then, by Proposition 9 (ii), $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$ for every integer $a \geq 3$. \square

By the discussion immediately preceding Proposition 4, it follows that for all positive integers a, b, c, d with $a \leq \min\{b, c, d\}$, the hypotheses of Proposition 10 are satisfied and hence we have proved Theorem 1.

4 Further properties of $F_{a,p,q,n}$ and $W_{a,p,q,n}$

Even if $p + q < 0$, there are still some situations in which $W_{a,p,q,n}(z)$ has only real roots. In this section we consider these, although they do not correspond to values of the parameters a, p, q and n that arise from rings of cliques. We need to apply the following result on the factorization of $F_{a,p,q,n}(z)$ when $a + p + n = 1$ or $a + p + n = 2$.

Proposition 11 Let a be an integer with $a \geq 1$ and p, q, n be arbitrary real numbers.

(i) If $a + p + n = 1$, then

$$F_{a,p,q,n}(z) = \prod_{j=0}^{a-1} (z^2 - (n+j)^2). \quad (23)$$

(ii) If $a + p + n = 2$, then

$$F_{a,p,q,n}(z) = (z^2 + (p-1)(n-1) + aq) \prod_{j=0}^{a-2} (z^2 - (n+j)^2). \quad (24)$$

Proof. (i) If $a + p + n = 1$, then

$$i!(a-i)! \binom{z-p}{a-i} \binom{z+n+i-1}{i} = \prod_{j=0}^{a-1} (z+n+j).$$

Thus

$$\begin{aligned} F_{a,p,q,n}(z) &= a! \sum_{i=0}^a i!(a-i)! \binom{a+p+q-1}{i} \binom{z-p}{a-i} \binom{z-q}{a-i} \binom{z+n+i-1}{i} \\ &= a! \prod_{j=0}^{a-1} (z+n+j) \sum_{i=0}^a \binom{a+p+q-1}{i} \binom{z-q}{a-i} \\ &= a! \prod_{j=0}^{a-1} (z+n+j) \binom{a+p+q-1+z-q}{a} \\ &= a! \prod_{j=0}^{a-1} (z+n+j) \binom{z-n}{a} \\ &= \prod_{j=0}^{a-1} (z^2 - (n+j)^2). \end{aligned} \quad (25)$$

Thus (i) holds.

(ii) Now let $a + p + n = 2$. Since $F_{1,p,q,n}(z) = z^2 + pq + pn + qn$, it is easy to verify that (ii) holds when $a = 1$. Assume that (ii) holds for any integer $1 \leq a < k$, where $k \geq 2$. Now let $a = k$.

Since $a + p + n = 2$, by Proposition 7,

$$F_{a,p,q,n}(z) = (z^2 + (a-1)(2p+2q+2n+2a-3) + pq + pn + qn) F_{a-1,p,q,n}(z).$$

As $a-1 + p + n = 1$, by (i) of this result, we have

$$F_{a-1,p,q,n}(z) = \prod_{j=0}^{a-2} (z^2 - (n+j)^2).$$

Since $p + n + a = 2$, it can be verified that

$$(a - 1)(2p + 2q + 2n + 2a - 3) + pq + pn + qn = (p - 1)(n - 1) + aq.$$

Hence (ii) also holds. \square

Proposition 12 *Let p, q, n be arbitrary real numbers with $p \leq q \leq n$.*

- (i) *If $p + q$ is a negative integer, then for every integer a with $a \geq 2 - p - q$, $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$.*
- (ii) *If $q + n$ is an integer, then for every integer a with $\max\{2, 2 - q - n\} \leq a \leq 2 - p - n$, $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$.*

Proof. (i) First consider the case that $a = 2 - p - q$. Since $p + q \leq -1$, we have $a \geq 3$. Proposition 11 implies that $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$.

Now assume that $a > 2 - p - q$ and $W_{a-1,p,q,n}(z)$ interleaves $W_{a-2,p,q,n}(z)$. Since $a > 2 - p - q$, we have $a + p + q - 2 \geq 1$ and so $a + q + n - 2 \geq a + p + n - 2 \geq a + p + q - 2 \geq 1$. Thus Proposition 9 (ii) implies that $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$. Therefore (i) holds.

(ii) The result is trivial if $\max\{2, 2 - q - n\} > 2 - p - n$. Now assume that $\max\{2, 2 - q - n\} \leq 2 - p - n$.

Let $a = \max\{2, 2 - q - n\}$. Then $a \geq 2 - q - n$, implying that $a + q + n - 2 \geq 0$. We also have $a \leq 2 - p - n$, implying that $a + p + n - 2 \leq 0$ and so $a + p + q - 2 \leq 0$. If $a = \max\{2, 2 - q - n\} = 2$, then Proposition 9 (i) implies that $W_{2,p,q,n}(z)$ interleaves $W_{1,p,q,n}(z)$, i.e., $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$. If $a = \max\{2, 2 - q - n\} = 2 - q - n$, then Proposition 11 implies that $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$.

Now assume that $\max\{2, 2 - q - n\} < a \leq 2 - p - n$ and $W_{a-1,p,q,n}(z)$ interleaves $W_{a-2,p,q,n}(z)$. Note that $\max\{2, 2 - q - n\} < a \leq 2 - p - n$ implies that $a + q + n - 2 > 0$ and $a + p + q - 2 \leq a + p + n - 2 \leq 0$. Thus Proposition 9 (ii) implies that $W_{a,p,q,n}(z)$ interleaves $W_{a-1,p,q,n}(z)$. Therefore (ii) holds. \square

By Proposition 12, the following result is obtained.

Proposition 13 *Let a be a positive integer and p, q, n be arbitrary real numbers with $p \leq q \leq n$. If one of the following conditions holds, then $W_{a,p,q,n}(z)$ has only real roots and therefore every root of $F_{a,p,q,n}(z)$ is either a real number or a complex number with its real part equal to 0:*

- (i) *$p + q$ is a negative integer and $a \geq 1 - p - q$; and*

(ii) $q + n$ is an integer and $\max\{1, 1 - q - n\} \leq a \leq 2 - p - n$. □

Acknowledgments

We wish to thank the Isaac Newton Institute for Mathematical Sciences, University of Cambridge, for generous support during the programme on Combinatorics and Statistical Mechanics (January-June 2008), where this work was started and partially finished.

References

- [1] Peter J. Cameron, Algebraic Properties of Chromatic Roots, *Available online at* www.maths.qmul.ac.uk/~pjc/csgnotes/alchrom1.pdf
- [2] Maria Chudnovsky and Paul Seymour, The roots of the independence polynomial of a clawfree graph, *J. Combin. Theory Ser. B* **97** (2007), 350–357.
- [3] F.M. Dong, K.M. Koh and K.L. Teo, *Chromatic polynomials and chromaticity of graphs*, World Scientific Publishing Co. Pte. Ltd., 2005.
- [4] Steve Fisk, *Polynomials, roots, and interlacing*, arXiv:math/0612833v2.
- [5] R.C. Read, An introduction to chromatic polynomials, *J. Combin. Theory* **4** (1968), 52–71.
- [6] R.C. Read, A large family of chromatic polynomials, in: *Proceedings of the Third Caribbean Conference on Combinatorics and Computing (Bridgetown, 1981)*, (1981), 23–41.
- [7] R.C. Read and W.T. Tutte, Chromatic polynomials, in: *Selected Topics in Graph Theory III* (eds. L.W. Beineke and R.J. Wilson), Academic Press, New York (1988), 15–42.
- [8] A.A. Zykov, On some properties of linear complexes, (Russian) *Mat. Sbornik N.S.* **24** (66) (1949), 163–188.